

# More clones from ideals

Martin Goldstern

Institute of Discrete Mathematics and Geometry  
Vienna University of Technology

Třešť, September 2008

# Outline

Background

Precomplete clones

Fixpoint clones

Ideal clones

Growth clones

# Outline

Background

Precomplete clones

Fixpoint clones

Ideal clones

Growth clones

## Definition

Fix a set  $X$ . We write  $\mathcal{O}^{(n)}$  for the set of  $n$ -ary operations:  
 $\mathcal{O}^{(n)} = X^{X^n}$ , and we let  $\mathcal{O} = \mathcal{O}_X = \bigcup_{n=1,2,\dots} \mathcal{O}^{(n)}$ .

A **clone on  $X$**  is a set  $C \subseteq \mathcal{O}$  which contains all the projection functions and is closed under composition.

Equivalently, a clone is the set of term functions of some universal algebra on  $X$ .

## Fact

The set of clones on  $X$  forms a complete Lattice: **CLONE**( $X$ ).

Definition: For any  $C \subseteq \mathcal{O}$  let  $\langle C \rangle$  be the clone generated by  $C$ .  
 We write  $C(f)$  for  $\langle C \cup \{f\} \rangle$ .

## Definition

Fix a set  $X$ . We write  $\mathcal{O}^{(n)}$  for the set of  $n$ -ary operations:  
 $\mathcal{O}^{(n)} = X^{X^n}$ , and we let  $\mathcal{O} = \mathcal{O}_X = \bigcup_{n=1,2,\dots} \mathcal{O}^{(n)}$ .

A **clone on  $X$**  is a set  $C \subseteq \mathcal{O}$  which contains all the projection functions and is closed under composition.

Equivalently, a clone is the set of term functions of some universal algebra on  $X$ .

## Fact

The set of clones on  $X$  forms a complete Lattice: **CLONE**( $X$ ).

Definition: For any  $C \subseteq \mathcal{O}$  let  $\langle C \rangle$  be the clone generated by  $C$ .  
 We write  $C(f)$  for  $\langle C \cup \{f\} \rangle$ .

## Definition

Fix a set  $X$ . We write  $\mathcal{O}^{(n)}$  for the set of  $n$ -ary operations:  
 $\mathcal{O}^{(n)} = X^{X^n}$ , and we let  $\mathcal{O} = \mathcal{O}_X = \bigcup_{n=1,2,\dots} \mathcal{O}^{(n)}$ .

A **clone on  $X$**  is a set  $C \subseteq \mathcal{O}$  which contains all the projection functions and is closed under composition.

Equivalently, a clone is the set of term functions of some universal algebra on  $X$ .

## Fact

The set of clones on  $X$  forms a complete Lattice: **CLONE**( $X$ ).

Definition: For any  $C \subseteq \mathcal{O}$  let  $\langle C \rangle$  be the clone generated by  $C$ .  
 We write  $C(f)$  for  $\langle C \cup \{f\} \rangle$ .

## Definition

Fix a set  $X$ . We write  $\mathcal{O}^{(n)}$  for the set of  $n$ -ary operations:  
 $\mathcal{O}^{(n)} = X^{X^n}$ , and we let  $\mathcal{O} = \mathcal{O}_X = \bigcup_{n=1,2,\dots} \mathcal{O}^{(n)}$ .

A **clone on  $X$**  is a set  $C \subseteq \mathcal{O}$  which contains all the projection functions and is closed under composition.

Equivalently, a clone is the set of term functions of some universal algebra on  $X$ .

## Fact

The set of clones on  $X$  forms a complete Lattice: **CLONE**( $X$ ).

Definition: For any  $C \subseteq \mathcal{O}$  let  $\langle C \rangle$  be the clone generated by  $C$ .  
 We write  $C(f)$  for  $\langle C \cup \{f\} \rangle$ .

## Size of $\mathbf{CLONE}(X)$

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then  $\mathbf{CLONE}(X)$  is countable, and completely understood. ("Post's Lattice")
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{|X|}$ , and not well understood.

If  $X$  is infinite, then

## Size of **CLONE**( $X$ )

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then **CLONE**( $X$ ) is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

## Size of $\mathbf{CLONE}(X)$

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then  $\mathbf{CLONE}(X)$  is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

## Size of **CLONE**( $X$ )

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then **CLONE**( $X$ ) is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then **CLONE**( $X$ ) =  $2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

- ▶  $|\mathcal{O}_X| = 2^{|X|}$ ,

## Size of **CLONE**( $X$ )

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then **CLONE**( $X$ ) is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

- ▶  $|\mathcal{O}_X| = 2^{|X|}$ ,
- ▶  $|\mathbf{CLONE}(X)| = 2^{2^{|X|}}$ ,
- ▶ and only little is known about the structure of **CLONE**( $X$ ).

## Size of **CLONE**( $X$ )

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then **CLONE**( $X$ ) is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

- ▶  $|\mathcal{O}_X| = 2^{|X|}$ ,
- ▶  $|\mathbf{CLONE}(X)| = 2^{2^{|X|}}$ ,
- ▶ and only little is known about the structure of **CLONE**( $X$ ).

## Size of **CLONE**( $X$ )

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then **CLONE**( $X$ ) is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

- ▶  $|\mathcal{O}_X| = 2^{|X|}$ ,
- ▶  $|\mathbf{CLONE}(X)| = 2^{2^{|X|}}$ ,
- ▶ and only little is known about the structure of **CLONE**( $X$ ).

## Size of $\mathbf{CLONE}(X)$

If  $X$  is finite, then  $\mathcal{O}_X$  is countable.

- ▶ If  $|X| = 1$ , then  $\mathcal{O}_X$  is trivial.
- ▶ If  $|X| = 2$ , then  $\mathbf{CLONE}(X)$  is countable, and completely understood. (“Post’s Lattice”)
- ▶ If  $3 \leq |X| < \aleph_0$ , then  $|\mathbf{CLONE}(X)| = 2^{\aleph_0}$ , and not well understood.

If  $X$  is infinite, then

- ▶  $|\mathcal{O}_X| = 2^{|X|}$ ,
- ▶  $|\mathbf{CLONE}(X)| = 2^{2^{|X|}}$ ,
- ▶ and only little is known about the structure of  $\mathbf{CLONE}(X)$ .

# Completeness

## Example

*The functions  $\wedge, \vee, \text{true}, \text{false}$  do not generate all operations on  $\{\text{true}, \text{false}\}$ .*

**Proof:** All these functions are monotone, and  $\neg$  is not.

Now let  $X$  be any set.

## Example

*Assume that  $\leq$  is a nontrivial partial order on  $X$ , and that all functions in  $C \subseteq \mathcal{O}$  are monotone with respect to  $\leq$ .*

*Then  $\langle C \rangle \neq \mathcal{O}$ .*

# Completeness

## Example

*The functions  $\wedge, \vee, \text{true}, \text{false}$  do not generate all operations on  $\{\text{true}, \text{false}\}$ .*

**Proof:** All these functions are monotone, and  $\neg$  is not.

Now let  $X$  be any set.

## Example

*Assume that  $\leq$  is a nontrivial partial order on  $X$ , and that all functions in  $C \subseteq \mathcal{O}$  are monotone with respect to  $\leq$ .*

*Then  $\langle C \rangle \neq \mathcal{O}$ .*

# Completeness

## Example

*The functions  $\wedge, \vee, \text{true}, \text{false}$  do not generate all operations on  $\{\text{true}, \text{false}\}$ .*

**Proof:** All these functions are monotone, and  $\neg$  is not.

Now let  $X$  be any set.

## Example

*Assume that  $\leq$  is a nontrivial partial order on  $X$ , and that all functions in  $C \subseteq \mathcal{O}$  are monotone with respect to  $\leq$ .*

*Then  $\langle C \rangle \neq \mathcal{O}$ .*

# Polymorphisms

Let  $X$  be a set,  $\mathcal{C} \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $\mathcal{C}$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $\mathcal{C}$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $\mathcal{C}$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subset A$ )

then  $\langle \mathcal{C} \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \emptyset$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \mathcal{O}$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Polymorphisms

Let  $X$  be a set,  $C \subseteq \mathcal{O}_X$ .

- ▶ If all functions in  $C$  respect some order  $\leq$  on  $X$ ,
- ▶ or: if all functions in  $C$  respect some nontrivial equivalence relation  $\theta$
- ▶ or: if all functions in  $C$  respect some nontrivial fixed set  $A \subset X$   
(i.e.,  $f[A^k] \subseteq A$ )
- ▶ or ...

then  $\langle C \rangle \neq \mathcal{O}$ .

We write  $\text{Pol}(\leq)$ ,  $\text{Pol}(\theta)$ ,  $\text{Pol}(A)$ , ... for the **clone of all functions respecting  $\leq$ ,  $\theta$ ,  $A$ , ...**

Instead of unary ( $A$ ) or binary ( $\leq$ ,  $\theta$ ) relations, we may also consider  $n$ -ary or even infinitary relations.

# Outline

Background

**Precomplete clones**

Fixpoint clones

Ideal clones

Growth clones

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(\theta_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “maximal” or “precomplete” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(A_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “maximal” or “precomplete” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

Question: Which relations  $R$  give rise to precomplete clones?

Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(A_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “maximal” or “precomplete” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(A_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “**maximal**” or “**precomplete**” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(\theta_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “**maximal**” or “**precomplete**” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

### Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(\theta_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “**maximal**” or “**precomplete**” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

### Question

*Which relations  $R$  give rise to precomplete clones?*

*This is nontrivial, already for binary relations.*

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(\theta_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “maximal” or “precomplete” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

### Question

*Which relations  $R$  give rise to precomplete clones?*

This is nontrivial, already for binary relations.

## Pol( ) and precomplete clones

- ▶ Every set of the form  $\text{Pol}(A_1) \cap \text{Pol}(A_2) \cap \text{Pol}(\theta_3) \cap \dots$  is a clone.
- ▶ Conversely, every clone is the intersection of sets of the form  $\text{Pol}(R)$  (where the  $R$ 's can be chosen of finite arity if  $X$  is finite).

The “maximal” or “precomplete” clones are the coatoms in the clone lattice.

$C \neq \emptyset$  is precomplete iff  $C(f) = \emptyset$  for all  $f \in \emptyset \setminus C$ .

- ▶ Every precomplete clone is of the form  $\text{Pol}(R)$  for some relation  $R$ .

### Question

*Which relations  $R$  give rise to precomplete clones?*

This is nontrivial, already for binary relations.

# Precomplete clones on finite sets

## Example

*Let  $\emptyset \neq A \neq X$ .*

*Then  $\text{Pol}(A)$  is precomplete.*

## Example

*Let  $X$  be finite. Let  $\theta$  be a nontrivial equivalence relation.*

*Then  $\text{Pol}(\theta)$  is precomplete.*

## Theorem (Rosenberg, 1970)

*There is an explicit list  $(R_i : i \in I)$  of finitely many (depending on the cardinality of  $X$ ) relations such that  $(\text{Pol}(R_i) : i \in I)$  lists all precomplete clones on  $X$ .*

*Moreover, every clone  $C \neq \emptyset$  is below some precomplete clone.*

# Precomplete clones on finite sets

## Example

*Let  $\emptyset \neq A \neq X$ .*

*Then  $\text{Pol}(A)$  is precomplete.*

## Example

*Let  $X$  be finite. Let  $\theta$  be a nontrivial equivalence relation.*

*Then  $\text{Pol}(\theta)$  is precomplete.*

## Theorem (Rosenberg, 1970)

*There is an explicit list  $(R_i : i \in I)$  of finitely many (depending on the cardinality of  $X$ ) relations such that  $(\text{Pol}(R_i) : i \in I)$  lists all precomplete clones on  $X$ .*

*Moreover, every clone  $C \neq \emptyset$  is below some precomplete clone.*

## Precomplete clones on finite sets

### Example

*Let  $\emptyset \neq A \neq X$ .*

*Then  $\text{Pol}(A)$  is precomplete.*

### Example

*Let  $X$  be finite. Let  $\theta$  be a nontrivial equivalence relation.*

*Then  $\text{Pol}(\theta)$  is precomplete.*

### Theorem (Rosenberg, 1970)

*There is an explicit list  $(R_i : i \in I)$  of finitely many (depending on the cardinality of  $X$ ) relations such that  $(\text{Pol}(R_i) : i \in I)$  lists all precomplete clones on  $X$ .*

*Moreover, every clone  $C \neq \emptyset$  is below some precomplete clone.*

## Precomplete clones on finite sets

### Example

*Let  $\emptyset \neq A \neq X$ .*

*Then  $\text{Pol}(A)$  is precomplete.*

### Example

*Let  $X$  be finite. Let  $\theta$  be a nontrivial equivalence relation.*

*Then  $\text{Pol}(\theta)$  is precomplete.*

### Theorem (Rosenberg, 1970)

*There is an explicit list  $(R_i : i \in I)$  of finitely many (depending on the cardinality of  $X$ ) relations such that  $(\text{Pol}(R_i) : i \in I)$  lists all precomplete clones on  $X$ .*

*Moreover, every clone  $C \neq \emptyset$  is below some precomplete clone.*

## Precomplete clones on finite sets

### Example

*Let  $\emptyset \neq A \neq X$ .*

*Then  $\text{Pol}(A)$  is precomplete.*

### Example

*Let  $X$  be finite. Let  $\theta$  be a nontrivial equivalence relation.*

*Then  $\text{Pol}(\theta)$  is precomplete.*

### Theorem (Rosenberg, 1970)

*There is an explicit list  $(R_i : i \in I)$  of finitely many (depending on the cardinality of  $X$ ) relations such that  $(\text{Pol}(R_i) : i \in I)$  lists all precomplete clones on  $X$ .*

Moreover, every clone  $C \neq \emptyset$  is below some precomplete clone.

# Precomplete clones on infinite sets

## Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

## Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

For which  $\Pi$  is  $\text{Pol}(\Pi)$  precomplete?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

► For which  $R$  is  $\text{Pol}(R)$  precomplete?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

► For which  $R$  is  $\text{Pol}(R)$  precomplete?

► Is every  $C \neq \emptyset$  below some precomplete clone?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

- ▶ For which  $R$  is  $\text{Pol}(R)$  precomplete?
- ▶ Is every  $C \neq \emptyset$  below some precomplete clone?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

- ▶ For which  $R$  is  $\text{Pol}(R)$  precomplete?
- ▶ Is every  $C \neq \emptyset$  below some precomplete clone?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

- ▶ For which  $R$  is  $\text{Pol}(R)$  precomplete?
- ▶ Is every  $C \neq \emptyset$  below some precomplete clone?

## Precomplete clones on infinite sets

### Example

Let  $\emptyset \neq A \neq X$ .

Then  $\text{Pol}(A)$  is precomplete.

### Example

Let  $\theta$  be a nontrivial equivalence relation **with finitely many classes**.

Then  $\text{Pol}(\theta)$  is precomplete.

- ▶ For which  $R$  is  $\text{Pol}(R)$  precomplete?
- ▶ Is every  $C \neq \emptyset$  below some precomplete clone?

# Outline

Background

Precomplete clones

**Fixpoint clones**

Ideal clones

Growth clones

# Fixpoint clones

## Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

## Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}(\langle F \rangle)$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,

$$\text{fix}(\langle F \rangle) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}.$$

$\text{fix}(\langle F \rangle)$  is a clone.

# Fixpoint clones

## Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

## Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

•  $\text{fix}((F))$  is a clone.

•  $\text{fix}((F))$  is the smallest clone generated by the set  $\{A \in F\}$ .

## Fixpoint clones

### Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

### Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

- ▶  $\text{fix}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{fix}((F)) = \text{fix}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.

# Fixpoint clones

## Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

## Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

- ▶  $\text{fix}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{fix}((F)) = \text{fix}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\Rightarrow$  maximal clone.

# Fixpoint clones

## Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

## Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

- ▶  $\text{fix}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{fix}((F)) = \text{fix}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\Rightarrow$  maximal clone.

## Fixpoint clones

### Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

### Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

- ▶  $\text{fix}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{fix}((F)) = \text{fix}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\Rightarrow$  maximal clone.

## Fixpoint clones

### Definition

Let  $A \subseteq X$ .  $\text{fix}(A)$  is the set of all functions  $f$  satisfying

$$\forall x \in A : f(x, \dots, x) = x.$$

This is a clone.

### Definition

Let  $F$  be a filter on  $X$ .  $\text{fix}((F))$  is defined as  $\bigcup_{A \in F} \text{fix}(A)$ , i.e.,  
 $\text{fix}((F)) = \{g : \exists A \in F \forall x \in A : g(x, \dots, x) = x\}$ .

- ▶  $\text{fix}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{fix}((F)) = \text{fix}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\Rightarrow$  maximal clone.

## Fixpoint clones, application

Let  $C_0 := \text{fix}(X)$ , i.e. the clone of all **idempotent** functions, i.e., functions  $f$  satisfying  $f(x, \dots, x) = x$  for *all*  $x \in X$ .

Let  $C_1 := \text{fix}(\emptyset) = \mathcal{O}$ , the clone of all functions. Then the interval  $[C_0, C_1]$  in the clone lattice is rather complicated, and yet we can “explicitly” describe it.

Theorem (Goldstern-Shelah, 2004)

*The clones in the interval  $[C_0, C_1]$  are exactly the clones  $\text{fix}(F)$ , for all possible filters (including the trivial filter  $\wp(X)$ ). (Maximal=precomplete clones correspond to ultrafilters.)*

So this interval is order isomorphic to the lattice of **closed subsets of  $\beta X$**  (with reverse inclusion).

## Fixpoint clones, application

Let  $C_0 := \text{fix}(X)$ , i.e. the clone of all **idempotent** functions, i.e., functions  $f$  satisfying  $f(x, \dots, x) = x$  for *all*  $x \in X$ .

Let  $C_1 := \text{fix}(\emptyset) = \mathcal{O}$ , the clone of all functions. Then the interval  $[C_0, C_1]$  in the clone lattice is rather complicated, and yet we can “explicitly” describe it.

Theorem (Goldstern-Shelah, 2004)

*The clones in the interval  $[C_0, C_1]$  are exactly the clones  $\text{fix}(F)$ , for all possible filters (including the trivial filter  $\mathfrak{P}(X)$ ). (Maximal=precomplete clones correspond to ultrafilters.)*

So this interval is order isomorphic to the lattice of **closed subsets of  $\beta X$**  (with reverse inclusion).

## Fixpoint clones, application

Let  $C_0 := \text{fix}(X)$ , i.e. the clone of all **idempotent** functions, i.e., functions  $f$  satisfying  $f(x, \dots, x) = x$  for *all*  $x \in X$ .

Let  $C_1 := \text{fix}(\emptyset) = \mathcal{O}$ , the clone of all functions. Then the interval  $[C_0, C_1]$  in the clone lattice is rather complicated, and yet we can “explicitly” describe it.

### Theorem (Goldstern-Shelah, 2004)

*The clones in the interval  $[C_0, C_1]$  are exactly the clones  $\text{fix}(F)$ , for all possible filters (including the trivial filter  $\wp(X)$ ). (Maximal=precomplete clones correspond to ultrafilters.)*

So this interval is order isomorphic to the lattice of **closed subsets of  $\beta X$**  (with reverse inclusion).

## Fixpoint clones, application

Let  $C_0 := \text{fix}(X)$ , i.e. the clone of all **idempotent** functions, i.e., functions  $f$  satisfying  $f(x, \dots, x) = x$  for *all*  $x \in X$ .

Let  $C_1 := \text{fix}(\emptyset) = \mathcal{O}$ , the clone of all functions. Then the interval  $[C_0, C_1]$  in the clone lattice is rather complicated, and yet we can “explicitly” describe it.

### Theorem (Goldstern-Shelah, 2004)

*The clones in the interval  $[C_0, C_1]$  are exactly the clones  $\text{fix}(F)$ , for all possible filters (including the trivial filter  $\wp(X)$ ). (Maximal=precomplete clones correspond to ultrafilters.)*

So this interval is order isomorphic to the lattice of **closed subsets of  $\beta X$**  (with reverse inclusion).

# Outline

Background

Precomplete clones

Fixpoint clones

**Ideal clones**

Growth clones

# Clones from ideals

## Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}(I)$  for the set of all functions preserving  $I$ .

►  $\text{Pol}(I)$  is a clone.

► If  $I$  is the principal ideal generated by the set  $A$ , then

$$\text{Pol}(I) = \text{Pol}(A).$$

► Every ideal is a large cube.

► Minimal ideal is maximal clone.

► Every maximal clone is a principal ideal.

► Every maximal clone is a large cube.

► Every maximal clone is a principal ideal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\Rightarrow$  larger clone.
- ▶  $\text{Pol}((I)) \supseteq \text{Pol}(A)$  for all  $A \in I$ .

# Clones from ideals

## Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\circ : [B]^\circ \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\circ : [B]^\circ \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Clones from ideals

### Definition

Let  $I$  be a nontrivial ideal on the set  $X$  containing all small sets.

$f : X^k \rightarrow X$  preserves  $I$  if  $\forall A \in I : f[A^k] \in I$ .

We write  $\text{Pol}((I))$  for the set of all functions preserving  $I$ .

- ▶  $\text{Pol}((I))$  is a clone.
- ▶ If  $I$  is the principal ideal generated by the set  $A$ , then  $\text{Pol}((I)) = \text{Pol}(A)$ .
- ▶ larger ideal  $\not\Rightarrow$  larger clone.
- ▶ maximal ideal  $\Rightarrow$  maximal clone.
- ▶ However, many other ideals also yield maximal clones.

$$I^\circ := \{A \subseteq X : \forall B \in [A]^\omega : [B]^\omega \cap I \neq \emptyset\}.$$

If  $I = I^\circ$ , then  $\text{Pol}((I))$  is maximal.

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  $2^{\aleph_0}$  many precomplete clones on a countable set. (Even without AC.)

Question

*Find such examples on uncountable sets.*

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  $2^c$  many precomplete clones on a countable set. (Even without AC.)

Question

*Find such examples on uncountable sets.*

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  $2^c$  many precomplete clones on a countable set. (Even without AC.)

Question

*Find such examples on uncountable sets.*

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

**Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)**

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  **$2^c$  many precomplete clones** on a countable set. (Even without AC.)

Question

*Find such examples on uncountable sets.*

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

**Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)**

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  $2^c$  many precomplete clones on a countable set. (Even without AC.)

### Question

*Find such examples on uncountable sets.*

## Ideal clones, application

For every subset  $A \subseteq 2^\omega$  we can find (explicitly) an ideal  $I_A$ , such that  $I_A = I_A^{-\circ}$ , and that the ideals  $I_A$  are all different.

**Theorem (Beiglböck-Goldstern-Heindorf-Pinsker, 2007)**

*While the ideals  $I_A$  are not maximal, the clones  $\text{Pol}((I_A))$  are (for nontrivial  $A$ ).*

This gives an explicit example of  $2^{\aleph_0}$  many precomplete clones on a countable set. (Even without AC.)

### Question

*Find such examples on uncountable sets.*

# Equivalence relations

## Example

*Let  $\theta$  be a nontrivial equivalence relation on a finite set. Then  $\text{Pol}(\theta)$  is a precomplete clone.*

## Example

*Let  $\theta$  be a nontrivial equivalence relation on any set, with finitely many classes. Then  $\text{Pol}(\theta)$  is a precomplete clone.*

## Equivalence relations

### Example

*Let  $\theta$  be a nontrivial equivalence relation on a finite set.  
Then  $\text{Pol}(\theta)$  is a precomplete clone.*

### Example

*Let  $\theta$  be a nontrivial equivalence relation on any set, with finitely many classes. Then  $\text{Pol}(\theta)$  is a precomplete clone.*

## Equivalence relations

### Example

*Let  $\theta$  be a nontrivial equivalence relation on a finite set. Then  $\text{Pol}(\theta)$  is a precomplete clone.*

### Example

*Let  $\theta$  be a nontrivial equivalence relation on any set, with finitely many classes. Then  $\text{Pol}(\theta)$  is a precomplete clone.*

## Definition

Let  $\mathcal{E}$  be a directed family of equivalence relations (coarser and coarser).

Define  $\text{Pol}(\mathcal{E})$  as the set of all functions  $f : X^k \rightarrow X$  with:  
for all  $E \in \mathcal{E}$  there is  $E' \in \mathcal{E}$  such that: whenever  $\vec{x} E \vec{y}$ , then  $f(\vec{x}) E' f(\vec{y})$ .

When is  $\text{Pol}(\mathcal{E})$  precomplete? Difficult. Because...

## Fact

For every ideal  $I$  there is a family  $\mathcal{E}$  as above such that  $\text{Pol}(I) = \text{Pol}(\mathcal{E})$ .

## Definition

Let  $\mathcal{E}$  be a directed family of equivalence relations (coarser and coarser).

Define  $\text{Pol}(\mathcal{E})$  as the set of all functions  $f : X^k \rightarrow X$  with:  
for all  $E \in \mathcal{E}$  there is  $E' \in \mathcal{E}$  such that: whenever  $\vec{x} E \vec{y}$ , then  $f(\vec{x}) E' f(\vec{y})$ .

When is  $\text{Pol}(\mathcal{E})$  precomplete? Difficult. Because...

## Fact

For every ideal  $I$  there is a family  $\mathcal{E}$  as above such that  $\text{Pol}(I) = \text{Pol}(\mathcal{E})$ .

## Definition

Let  $\mathcal{E}$  be a directed family of equivalence relations (coarser and coarser).

Define  $\text{Pol}(\mathcal{E})$  as the set of all functions  $f : X^k \rightarrow X$  with:  
for all  $E \in \mathcal{E}$  there is  $E' \in \mathcal{E}$  such that: whenever  $\vec{x} E \vec{y}$ , then  $f(\vec{x}) E' f(\vec{y})$ .

When is  $\text{Pol}(\mathcal{E})$  precomplete? Difficult. Because...

## Fact

For every ideal  $I$  there is a family  $\mathcal{E}$  as above such that  $\text{Pol}(I) = \text{Pol}(\mathcal{E})$ .

## Definition

Let  $\mathcal{E}$  be a directed family of equivalence relations (coarser and coarser).

Define  $\text{Pol}(\mathcal{E})$  as the set of all functions  $f : X^k \rightarrow X$  with:  
for all  $E \in \mathcal{E}$  there is  $E' \in \mathcal{E}$  such that: whenever  $\vec{x} E \vec{y}$ , then  $f(\vec{x}) E' f(\vec{y})$ .

When is  $\text{Pol}(\mathcal{E})$  precomplete? Difficult. Because...

## Fact

For every ideal  $I$  there is a family  $\mathcal{E}$  as above such that  $\text{Pol}(I) = \text{Pol}(\mathcal{E})$ .

# Outline

Background

Precomplete clones

Fixpoint clones

Ideal clones

**Growth clones**

# Growth clones

## Definition

Let  $X = \mathbb{N} = \{0, 1, 2, \dots\}$  for simplicity. For every infinite  $A = \{a_0 < a_1 < \dots\} \subseteq X$  we define **bound**( $A$ ) as the set of functions which do not jump to far in  $A$ :

$$\text{bound}(A) := \{f : \exists k \forall i : \vec{x} < a_i \Rightarrow f(\vec{x}) < a_{i+k}\}$$

*(This is a clone.)*

A similar construction is possible for uncountable sets.

# Growth clones

## Definition

Let  $X = \mathbb{N} = \{0, 1, 2, \dots\}$  for simplicity. For every infinite  $A = \{a_0 < a_1 < \dots\} \subseteq X$  we define **bound**( $A$ ) as the set of functions which do not jump to far in  $A$ :

$$\text{bound}(A) := \{f : \exists k \forall i : \vec{x} < a_i \Rightarrow f(\vec{x}) < a_{i+k}\}$$

*(This is a clone.)*

A similar construction is possible for uncountable sets.

# Growth clones

## Definition

Let  $X = \mathbb{N} = \{0, 1, 2, \dots\}$  for simplicity. For every infinite  $A = \{a_0 < a_1 < \dots\} \subseteq X$  we define **bound**( $A$ ) as the set of functions which do not jump to far in  $A$ :

$$\text{bound}(A) := \{f : \exists k \forall i : \vec{x} < a_i \Rightarrow f(\vec{x}) < a_{i+k}\}$$

*(This is a clone.)*

A similar construction is possible for uncountable sets.

# Growth clones

## Definition

Let  $X = \mathbb{N} = \{0, 1, 2, \dots\}$  for simplicity. For every infinite  $A = \{a_0 < a_1 < \dots\} \subseteq X$  we define **bound**( $A$ ) as the set of functions which do not jump to far in  $A$ :

$$\text{bound}(A) := \{f : \exists k \forall i : \vec{x} < a_i \Rightarrow f(\vec{x}) < a_{i+k}\}$$

*(This is a clone.)*

A similar construction is possible for uncountable sets.

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\not\Rightarrow$  maximal clone.
- ▶ (In fact,  $\text{bound}((F))$  is never a maximal clone.)

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\not\Rightarrow$  maximal clone.
- ▶ (In fact,  $\text{bound}((F))$  is never a maximal clone.)

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\not\Rightarrow$  maximal clone.
- ▶ (In fact,  $\text{bound}((F))$  is never a maximal clone.)

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\not\Rightarrow$  maximal clone.
- ▶ (In fact,  $\text{bound}((F))$  is never a maximal clone.)

## Growth clones, continued

### Definition

Let  $X = \mathbb{N}$  again. For every filter  $F$  of subsets of  $X$  we define  $\text{bound}((F)) := \bigcup_{A \in F} \text{bound}(A)$ .

$$\text{bound}((F)) := \{f : \exists A \in F \exists k \forall i : \vec{x} < a_i^A \Rightarrow f(\vec{x}) < a_{i+k}^A\}$$

(where  $a_0^A < a_1^A < \dots$  is the increasing enumeration of  $A$ ).

- ▶  $\text{bound}((F))$  is a clone.
- ▶ If  $F$  is the principal filter generated by the set  $A$ , then  $\text{bound}((F)) = \text{bound}(A)$ .
- ▶ larger filter  $\Rightarrow$  larger clone.
- ▶ maximal filter  $\not\Rightarrow$  maximal clone.
- ▶ (In fact,  $\text{bound}((F))$  is never a maximal clone.)

## Growth clones, application

Theorem ( $G^*$ -Shelah, 2002)

Assume CH. Then on there is a filter  $F$  on  $\mathbb{N} = \{0, 1, 2, \dots\}$  such that, letting  $C := \text{bound}((F))$ , we know the interval  $[C, \mathcal{O})$  quite well: it is (more or less) a quite saturated linear order  $\mathbf{L}$  with no last element.

(In particular: not every clone is below a precomplete clone.)

We can choose  $\text{bound}((F))$  in such a way that the relation  $f \leq g \Leftrightarrow f \in C(g)$  is a linear quasiorder. The clones above  $C$  will then be the Dedekind cuts in this order.

This relation  $f \leq g$  means that on a large set (i.e., a set in the filter  $F$ ),  $g$  grows at least as fast as  $f$ .

## Growth clones, application

### Theorem ( $G^*$ -Shelah, 2002)

*Assume CH. Then on there is a filter  $F$  on  $\mathbb{N} = \{0, 1, 2, \dots\}$  such that, letting  $C := \text{bound}((F))$ , we know the interval  $[C, \mathcal{O})$  quite well: it is (more or less) a quite saturated linear order  $\mathbf{L}$  with no last element.*

*(In particular: not every clone is below a precomplete clone.)*

We can choose  $\text{bound}((F))$  in such a way that the relation  $f \leq g \Leftrightarrow f \in C(g)$  is a linear quasiorder. The clones above  $C$  will then be the Dedekind cuts in this order.

This relation  $f \leq g$  means that on a large set (i.e., a set in the filter  $F$ ),  $g$  grows at least as fast as  $f$ .

## Growth clones, application

### Theorem ( $G^*$ -Shelah, 2002)

Assume CH. Then on there is a filter  $F$  on  $\mathbb{N} = \{0, 1, 2, \dots\}$  such that, letting  $C := \text{bound}((F))$ , we know the interval  $[C, \infty)$  quite well: it is (more or less) a quite saturated linear order  $\mathbf{L}$  with no last element.

(In particular: not every clone is below a precomplete clone.)

We can choose  $\text{bound}((F))$  in such a way that the relation  $f \leq g \Leftrightarrow f \in C(g)$  is a linear quasiorder. The clones above  $C$  will then be the Dedekind cuts in this order.

This relation  $f \leq g$  means that on a large set (i.e., a set in the filter  $F$ ),  $g$  grows at least as fast as  $f$ .

## Growth clones, new application

Theorem? (Aug-Sep 2008)

Let  $\mathbb{N} = N_1 \dot{\cup} N_2$ , with two infinite disjoint sets  $N_1, N_2$ , say odd and even numbers.

Assume CH. Then there are filters  $F_1, F_2$  on  $N_1$  and  $N_2$ , respectively, such that, letting  $C := \text{bound}((F_1)) \cap \text{bound}((F_2))$ , we know the interval  $[C, \emptyset)$  quite well: it is (more or less)  $\mathbf{L} \times \mathbf{L}$ , with  $\mathbf{L}$  the quite saturated linear order from the previous slide.

Theorem? (2009?)

Let  $(F_i : i \in I)$  be a family of many (almost?) disjoint sets.

Assume CH. Then there filters  $F_i, i \in I$ , such that, letting  $C := \bigcap_{i \in I} \text{bound}((F_i))$ , the interval  $[C, \emptyset)$  is (more or less)  $\mathbf{L}^I$ .

## Growth clones, new application

### Theorem? (Aug-Sep 2008)

Let  $\mathbb{N} = N_1 \dot{\cup} N_2$ , with two infinite disjoint sets  $N_1, N_2$ , say odd and even numbers.

Assume CH. Then there are filters  $F_1, F_2$  on  $N_1$  and  $N_2$ , respectively, such that, letting  $C := \text{bound}((F_1)) \cap \text{bound}((F_2))$ , we know the interval  $[C, \emptyset)$  quite well: it is (more or less)  $\mathbf{L} \times \mathbf{L}$ , with  $\mathbf{L}$  the quite saturated linear order from the previous slide.

### Theorem? (2009?)

Let  $(F_i : i \in I)$  be a family of many (almost?) disjoint sets.

Assume CH. Then there filters  $F_i, i \in I$ , such that, letting  $C := \bigcap_{i \in I} \text{bound}((F_i))$ , the interval  $[C, \emptyset)$  is (more or less)  $\mathbf{L}^I$ .

## Growth clones, new application

### Theorem? (Aug-Sep 2008)

Let  $\mathbb{N} = N_1 \dot{\cup} N_2$ , with two infinite disjoint sets  $N_1, N_2$ , say odd and even numbers.

Assume CH. Then there are filters  $F_1, F_2$  on  $N_1$  and  $N_2$ , respectively, such that, letting  $C := \text{bound}((F_1)) \cap \text{bound}((F_2))$ , we know the interval  $[C, \emptyset)$  quite well: it is (more or less)  $\mathbf{L} \times \mathbf{L}$ , with  $\mathbf{L}$  the quite saturated linear order from the previous slide.

### Theorem? (2009?)

Let  $(F_i : i \in I)$  be a family of many (almost?) disjoint sets.

Assume CH. Then there filters  $F_i, i \in I$ , such that, letting  $C := \bigcap_{i \in I} \text{bound}((F_i))$ , the interval  $[C, \emptyset)$  is (more or less)  $\mathbf{L}^I$ .